

Heavy Quark Productions with Spin*

Toshiyuki Mori

Faculty of Human Development, Kobe University, Nada, Kobe 657, Japan

Abstract

Heavy quark productions in high energy polarized scatterings are reviewed from a personal point of view. After mentioning why heavy quark physics is so interesting, I concentrate on two rather specific subjects: (1) polarized Λ_c^+ productions in deep inelastic ℓp scatterings and (2) ψ' productions in polarized pp collisions. The first one is an interesting process for extracting the polarized gluon in the proton and the second one may give another test of the color-octet model of heavy quarkonium productions in high energy collisions.

1. Why heavy quarks?

Quantum Chromodynamics(QCD) is the underlying field theory of strong interactions of quarks and gluons. Although it is surprisingly successful in describing physics in perturbative regions, it is not so easy task to directly apply QCD for nonperturbative regions because of its complicated structure. To understand physics of quarks and gluons, it is not enough to write down the QCD Lagrangian. The more important thing is to study how quarks and gluons interact to make hadrons and how they are produced in high energy collisions.

So far, to understand the complicated structure of hadron dynamics in nonperturbative regions, many effective theories[1] such as the potential model, the chiral perturbation theory, the Skyrme model, the Bag model, the heavy quark effective theory(HQET) and so on, have been proposed. These theories have their own scales in which they work. Here I would like to just mention an important role of mass hierarchy and scale. At present, we have 6 quark flavors, u , d , s , c , b , and t , whose masses are arranged, in order, from light to heavy quarks. Among them, the masses of u , d and s quarks are smaller than the QCD scale $\Lambda_{QCD}(\approx 200MeV)$ and the SU(3) flavor symmetry is approximately realized as a good symmetry[2]. Chiral dynamics works well for these light quarks and can be applied for parameters remaining to be constant when $m_q \rightarrow 0$. Actually m_q is not 0 and we have symmetry breaking due to m_q/Λ_{QCD} . On the other hand, the heavy quark effective theory(HQET)[3] works well for the c and b quarks which are much heavier than the Λ_{QCD} value. For hadrons containing these quarks, the heavy quark spin interaction decouples from QCD interactions and thus the SU($2N_f$) spin-flavor symmetry appears as a good symmetry. The HQET can be applied for parameters remaining to be constant when $m_Q \rightarrow \infty$. Since m_Q is actually not infinite, we have symmetry breaking due to Λ_{QCD}/m_Q . Practically, the HQET can not be applied for hadrons including a top quark. The top quark is very special because it is extremely heavy: it is much heavier than the W boson with $m_W \approx 80GeV$. A top quark decays into a b quark emitting a W boson as

*Talk given at the International School Seminar'97, "Structure of Particles and Nuclei and their Interactions", (Tashkent, Oct. 6-13, 1997)

a real process and a single top decay width becomes very large, $\Gamma(t \rightarrow bW) \approx 1.5\text{GeV}$ for $m_t = 175\text{GeV}$. Therefore, the life time of the t quark becomes $\approx 10^{-25}\text{sec}$ which is shorter than the hadronization time $\approx 10^{-23}\text{sec}$ and there is no possibility to make hadrons containing top quarks[4].

In summary, physics of the u , d and s quarks is something similar to the solid state physics. Both of them are described well by effective theories with beautiful symmetries. On the other hand, since a top quark is too heavy to make hadrons containing it, we do not need to worry about the complicated nonperturbative effect of its hadronization. One can directly test the perturbative QCD in top physics. The situation is, in some sense, similar to a gas: both of them are rather simple systems. However, physics of charm and bottom quarks is far from these two limits. It is something similar to physics of liquid which is in between solid and gas. To understand physics of charm and bottom quarks, we need knowledge of both perturbative and nonperturbative QCD. In other words, heavy quarks, i.e. charm and bottom quarks provide an important playground for testing both perturbative and nonperturbative QCD. For the nonperturbative and static region, the potential model approach is still effective in the spectroscopy of these heavy flavored hadrons in addition to the HQET[5]. On the other hand, the heavy quarks are produced only via gluon interactions in high energy scatterings and thus play an important role in extracting an information on perturbative QCD.

Recently, there have been new interests in physics of heavy flavored quarks: how they work effectively for extracting the information on the spin structure of nucleons and also how they are produced in high energy collisions. In this talk, I concentrate on two topics related to these subjects which we have recently studied.

2. Polarized structure functions of nucleons

Recent high energy polarized experiments have revealed a much more fruitful structure of nucleons than ever considered. One of the most serious problems is so-called the spin puzzle[6], i.e.

$$\frac{1}{2} = \frac{1}{2}\Delta\Sigma + \Delta g + \langle L_Z \rangle_q + \langle L_Z \rangle_g, \quad (1)$$

$$\Delta\Sigma = \Delta u + \Delta d + \Delta s \approx 0.3, \quad (2)$$

$$\Delta s \approx -0.12, \quad (3)$$

where $\Delta\Sigma$ and Δg are the amount of the proton spin carried by quarks and gluons, respectively, and $\langle L_Z \rangle_{q,g}$ implies the orbital angular momenta of quarks and gluons. Although there have been many theoretical and experimental studies so far, many questions are still to be answered: where does the proton spin come from?, why are s quarks polarized negatively?, what about gluons?, how does QCD work? and so on.

In order to go beyond the present understanding on the nucleon spin structure, it is very important to measure the polarized gluon and sea-quark distribution. Here I am interested in the gluon polarization in the proton. Although there have been many discussions on the gluon polarization, knowledge of the magnitude Δg and the behavior $\delta g(x, Q^2)$ is still poor. Among many processes which are sensitive to δg , here we propose a different process(fig.1),




Figure 1: The lowest order QCD diagram for Λ_c^+ lepton productions in unpolarized lepton-polarized proton scatterings.

$$\ell + \vec{p} \rightarrow \vec{\Lambda}_c^+ + X, \quad (4)$$

to get further information on polarized gluons, expecting that the process can be observed in the forthcoming COMPASS experiment, where the arrow attached to particles means that these particles are polarized. The process is expected to be effective for testing δg since its cross section is directly proportional to δg . Furthermore, the spin of Λ_c^+ is carried by the c quark and thus measurement of polarization of Λ_c^+ in the target fragmentation region could determine the gluon polarization, δg .

An interesting parameter is the two-spin asymmetry,

$$A_{LL} = \frac{[d\sigma_{++} - d\sigma_{+-} + d\sigma_{--} - d\sigma_{-+}]}{[d\sigma_{++} + d\sigma_{+-} + d\sigma_{--} + d\sigma_{-+}]} = \frac{d\Delta\sigma/dy}{d\sigma/dy}, \quad (5)$$

where $d\sigma_{+-}$, for instance, denotes that the spin of the target proton and produced Λ_c^+ is positive and negative, respectively. The explicit expressions of the spin-independent and spin-dependent cross sections are given in ref.[7]. Using the typical examples of polarized gluon distributions(GS95[8], BBS95[9], GRV95[10])(fig.2), we have calculated the A_{LL} at a CMS energy of the virtual photon-proton collision, $\sqrt{s} = 10\text{GeV}$ (which corresponds to γ^* energy $\nu = 56\text{GeV}$) and a momentum transfer squared $Q^2 = 10\text{GeV}^2$ (fig.3), whose kinematical region can be covered by COMPASS experiments. We have found that the A_{LL} largely depends on the behavior of polarized gluons. Thus we can say that the process might be promising to test various models of polarized gluons.




Figure 2: The x -dependence of polarized gluon distributions at $Q^2=10\text{GeV}^2$. The solid, dotted, dash-dotted and dashed lines indicate the set A, C of ref.[8], ref.[9] and the 'standard scenario' of ref.[10], respectively.




Figure 3: The spin correlation asymmetry A_{LL} as a function of rapidity y at $\sqrt{s}=10\text{GeV}$ and $Q^2=10\text{GeV}^2$. Various lines represent the same as in fig.2.

3. Heavy quarkonium productions

So far, the standard model is extremely successful in describing the present experimental data. However, recent observation[11] by the CDF collaboration has shown that the production cross section of charmonium at large p_T region in pp collisions at Tevatron are order of magnitude larger than the conventional QCD prediction by the color-singlet model. This dramatic discrepancy might make a new step toward the deep understandings of heavy quarkonium production mechanism. To remove such a big discrepancy between the experimental data and the prediction of the color-singlet model, an interesting new color-octet model has been proposed recently by several people[12]. Physics of the color-octet model whose theoretical ground has been rigorously formulated by a new effective theory called the nonrelativistic QCD(NRQCD)[13], is now one of the most interesting topics for heavy quarkonium productions at high energy. Although several processes have been already suggested for testing the model[14], the discussion seems still controvertial[15]. To go beyond the present theoretical understandings, it is necessary to study other processes. To test the model, we propose here another process,

$$\vec{p} + \vec{p} \rightarrow \psi' + X, \quad (6)$$

at small p_T regions[16]. Since the process is dominated by the gluon-gluon fusion, there is no direct production of color-singlet ψ' because of charge conjugation. There are only two states which are expected to contribute to the ψ' production in the final states: the color-octet state decaying to $\psi' + g$ and the 2^3P_2 state decaying to $\psi' + \gamma$. By using typical examples of polarized gluon distributions presented in fig.2, we have calculated the two-spin asymmetry A_{LL} for the ψ' produced in this process and found that it is positive for the color-octet state and negative for the 2^3P_2 state. The spin-dependent and

spin-independent cross sections of ψ' productions via the color-octet state depend on the parameters, $\langle \mathcal{O}_8^{\psi'}(^1S_0) \rangle$, $\langle \mathcal{O}_8^{\psi'}(^3S_1) \rangle$ and $\langle \mathcal{O}_8^{\psi'}(^3P_0) \rangle$ [17], which are nonperturbative long-distance factors associated with the production of a $c\bar{c}$ pair in a color-octet 1S_0 , 3S_1 and 3P_0 states, respectively. The model can also be applied likewise even for the Υ' production. These nonperturbative factors are obtained from recent analysis on charmonium and bottomonium hadroproductions:

$$\langle \mathcal{O}_8^{\psi'}(^3S_1) \rangle \approx 4.6 \times 10^{-3} [\text{GeV}^3], \quad (7)$$

$$\langle \mathcal{O}_8^{\psi'}(^1S_0) \rangle + \frac{7}{m_c^2} \langle \mathcal{O}_8^{\psi'}(^3P_0) \rangle \approx 5.2 \times 10^{-3} [\text{GeV}^3], \quad (8)$$

$$\frac{1}{3} \langle \mathcal{O}_8^{\psi'}(^1S_0) \rangle + \frac{1}{m_c^2} \langle \mathcal{O}_8^{\psi'}(^3P_0) \rangle \approx (5.9 \pm 1.9) \times 10^{-3} [\text{GeV}^3], \quad (9)$$

for ψ' productions[18] and

$$\langle \mathcal{O}_8^{\Upsilon'}(^3S_1) \rangle \approx 4.1 \times 10^{-3} [\text{GeV}^3], \quad (10)$$

$$\langle \mathcal{O}_8^{\Upsilon'}(^1S_0) \rangle + \frac{7}{m_b^2} \langle \mathcal{O}_8^{\Upsilon'}(^3P_0) \rangle \approx 3.0 \times 10^{-2} [\text{GeV}^3], \quad (11)$$

$$\frac{1}{5} \langle \mathcal{O}_8^{\Upsilon'}(^1S_0) \rangle + \frac{1}{m_b^2} \langle \mathcal{O}_8^{\Upsilon'}(^3P_0) \rangle \approx (9.1 \pm 7.2) \times 10^{-3} [\text{GeV}^3], \quad (12)$$

for Υ' productions[18][19]. The A_{LL} via the color-octet state largely depend on the ratio, $\frac{\bar{\Theta}}{\Theta} \equiv \frac{\langle \mathcal{O}_8(^1S_0) \rangle - \frac{1}{m^2} \langle \mathcal{O}_8(^3P_0) \rangle}{\langle \mathcal{O}_8(^1S_0) \rangle + \frac{7}{m^2} \langle \mathcal{O}_8(^3P_0) \rangle}$, whose values range as $\frac{\bar{\Theta}}{\Theta} \approx 3.6 - 8.0$ for ψ' productions and $\frac{\bar{\Theta}}{\Theta} \approx -1.88 - 8.01$ for Υ' productions. Those for the 2^3P_2 state depend on the derivative of the wave function at the origin, $|R'_{2^3P_2}(0)|$, whose value has been estimated by the potential models[20]; $|R'_{2^3P_2}(0)| = 0.076\text{GeV}^5 - 0.186\text{GeV}^5$ for ψ' and $|R'_{2^3P_2}(0)| = 1.417\text{GeV}^5 - 2.067\text{GeV}^5$ for Υ' , depending on the type of potentials. We have found that the A_{LL} for the sum of contributions from the color-octet state and the 2^3P_2 state becomes positive for the present parameter regions, in particular, at $\sqrt{s} = 50\text{GeV}$: the results with typical parameters are shown in figs.4 and 5. From this result, one can conclude that if we observe a positive A_{LL} in the future RHIC experiment, we can definitely say that the color-octet model really contribute to this process. The process is therefore very effective for testing the color-octet model. Since the results largely depend on the ratio, $\frac{\bar{\Theta}}{\Theta}$, it is very important to determine the value from other experiments in order to give a better prediction. Furthermore, the process is effective for testing the spin-dependent gluon distribution in the proton because its cross section is directly proportional to the product of $\delta g(x)$ in both protons.

4. Outlook

The production of heavy quarks and quarkonia is a very important subject to test the standard model and QCD, and furthermore to go beyond the present understandings of particle physics. Polarized experiments are the most promising way for testing various theories on heavy quark physics with spin. In addition to the running experiments, a lot of interesting polarized experiments, such as, COMPASS experiment at CERN, RHIC project, future polarized HERA experiments, polarized γp experiments at SPring-8(Japan), MAMI(Germany), TJNAF(USA), and so on, are planned at many places in the




Figure 4: The two-spin asymmetry $A_{LL}^{\psi'}(pp)$ for $\frac{\tilde{\Theta}}{\Theta} = 5.8$, $|R'_{2^3P_2}(0)|^2 = 0.131\text{GeV}^5$ at $\sqrt{s} = 50\text{GeV}$, calculated with various types of $\Delta g(x)$, as a function of longitudinal momentum fraction x_L of ψ' . Various lines are indicated in the figure.




Figure 5: The two-spin asymmetry $A_{LL}^{\Upsilon'}(pp)$ for $\frac{\tilde{\Theta}}{\Theta} = 3.06$, $|R'_{2^3P_2}(0)|^2 = 1.572\text{GeV}^5$ at $\sqrt{s} = 50\text{GeV}$, calculated with various types of $\Delta g(x)$, as a function of longitudinal momentum fraction x_L of Υ' . Various lines are indicated in the figure.

world and will come out soon. These experiments will lead us to a new stage for studying heavy quark physics with spin.

While a lot of experimental and theoretical progress has been done so far for the spin physics and heavy quark productions, heavy quark physics is still challenging subject which needs further investigation.

References

- [1] J. H. Donoghue, E. Golowich and B. R. Holstein, "Dynamics of the Standars Model", Cambridge university press (1992).
- [2] H. Georgi, "Weak Interactions and Modern Particle Theory", Benjamin/Cummings Pub., (1984).
- [3] M. Neubert, Phys. Rep. **245**, 259 (1994).
- [4] I. Bigi, Y. Dokshitzer, V. Khoze, J. H. Kühn and P. M. Zerwas, Phys. Lett. **B181**, 151 (1986).
- [5] T. Matsuki and T. Morii, Phys. Rev. **D56**, 5646 (1997).
- [6] H. -Y. Cheng, hep-ph/9607254 (1996).
- [7] N. Kochelev, T. Morii and T. Yamanishi, Phys. Lett. **B405**, 168 (1997).
- [8] T. Gehrmann and W. J. Stirling, Z. Phys. **C65**, 461 (1995).

- [9] S. J. Brodsky, M. Burkardt and I. Schmidt, Nucl. Phys. **B441**, 197 (1995).
- [10] M. Glück, E. Reya and W. Vogelsang, Phys. Lett. **B359**, 201 (1995).
- [11] F. Abe et. al., CDF Collab., Phys. Rev. Lett. **69** 3704 (1992); Phys. Rev. Lett. **71** 2537 (1993).
- [12] E. Braaten and S. Fleming, Phys. Rev. Lett. **74** 3327 (1995); P. Cho and A. K. Leibovich, Phys. Rev. **D53** 150 (1996); M. Cacciari, M. Greco, M.L. Mangano and A. Petrelli, Phys. Lett. **B356** 553 (1995).
- [13] G. T. Bodwin, E. Braaten and G. P. Lepage, Phys. Rev. **D51** 1125 (1995).
- [14] P. Cho and A. K. Leibovich, Phys. Rev. **D53** 6203 (1996); E. Braaten and Y. -Q. Chen, Phys. Rev. Lett. **76** 730 (1996); K. Cheung, W. -Y. Keung and T. C. Yuan, Phys. Rev. Lett. **76** 877 (1996); P. Cho, Phys. Lett. **B368** 171 (1996).
- [15] M. Cacciari and M. Krämer, Phys. Rev. Lett. **76** 4128 (1996).
- [16] T. Morii, D. Roy, K. Sudoh, Zuxun Sun, S. Tanaka and T. Yamanishi, hep-ph/9708415 (1997).
- [17] S. Gupta and P. Mathews, preprint TIFR/TH/96-53, hep-ph/9609504 (1996).
- [18] M. Beneke and I. Z. Rothstein, Phys. Rev. **D54** 2005 (1996); A. K. Leibovich, preprint CALT-68-2082, hep-ph/9610381 (1996).
- [19] P. Cho and A. K. Leibovich, Phys. Rev. **D53** 6203 (1996).
- [20] E. J. Eichten and C. Quigg, Phys. Rev. **D52** 1726 (1995).